

ZIKA & PREGNANCY



Please visit the Center for Disease Control website for the most current updates and travel alert
www.cdc.gov/zika

Common-sense approach to avoid birth defects linked to ZIKA:
It's about having a healthy BABY

- **AVOID travel to Zika affected areas if you are of child-birth age**
- Do not have unprotected sex with exposed males who have recently traveled to Zika affected areas
- **Prevention depends on the behavior of the prospective parent**
- **Couples who want to conceive:**
 - Asymptomatic men and women should wait 8 weeks after travel to Zika affected areas
 - Symptomatic women should wait 8 weeks after recovery
 - Symptomatic men should wait 6 months after recovery
- **There is no current vaccine to prevent Zika or medicine to treat Zika**



PREGNANT and living in an area with Zika?



What we know about Zika

- ◆ Zika can be passed from a pregnant woman to her fetus.
- ◆ Infection with Zika during pregnancy is linked to birth defects in babies.
- ◆ Zika is spread mostly by the bite of an infected *Aedes* species mosquito.
 - » These mosquitoes are aggressive daytime biters. They also bite at night.
- ◆ There is no vaccine to prevent or medicine to treat Zika.
- ◆ Zika can be spread by a man to his sex partners.



What we don't know about Zika


- ◆ When during pregnancy Zika might cause harm to the fetus.
- ◆ How likely it is that Zika infection will affect your pregnancy.
- ◆ If your baby will develop birth defects from the infection.

Symptoms of Zika

Most people with Zika won't even know they have it. The illness is usually mild with symptoms lasting for several days to a week.

The most common symptoms of Zika are

- ◆ Fever
- ◆ Rash
- ◆ Joint Pain
- ◆ Red eyes



U.S. Department of Health and Human Services
Centers for Disease Control and Prevention

CDC recommends special precautions for pregnant women and women trying to become pregnant

Pregnant?

Pregnant women and their male partners should strictly follow steps to prevent mosquito bites.

If you have a male partner, either use condoms the right way every time you have sex during your pregnancy, or do not have sex.

If you develop the symptoms of Zika, see a healthcare provider right away for testing.

Testing is recommended for pregnant women at their first prenatal care visit.



Trying to become pregnant?

Women trying to become pregnant and their male partners should strictly follow steps to prevent mosquito bites.

Talk to your healthcare provider about plans to become pregnant.

Your Best Protection: Prevent Mosquito Bites

Clothing

- ◆ Wear long-sleeved shirts and long pants.
- ◆ Treat clothing and gear with permethrin or purchase permethrin-treated items.
 - » Treated clothing remains protective after multiple washings. See product information to learn how long the protection will last.
 - » If treating items yourself, follow the product instructions carefully.
- ◆ Do NOT use permethrin products directly on skin. They are intended to treat clothing.



Indoor Protection

- ◆ Stay in places with air conditioning or that use window and door screens to keep mosquitoes outside.
- ◆ Sleep under a mosquito bed net if air conditioned or screened rooms are not available or if sleeping outdoors.



Repellent

Use Environmental Protection Agency (EPA)-registered insect repellents. When used as directed, these insect repellents are safe and effective for pregnant and breastfeeding women.

- ◆ Always follow the product label instructions.
- ◆ Reapply as directed.
- ◆ Do not spray repellent on the skin under clothing.
- ◆ If you are also using sunscreen, apply sunscreen before applying insect repellent.



ZIKA & PREGNANCY



Please visit the Center for Disease Control website for the most current updates and travel alert
www.cdc.gov/zika